

The group G is isomorphic to the group labelled by [12, 1] in the Small Groups library.

Ordinary character table of $G \cong \text{C3} : \text{C4}$:

	1a	2a	3a	6a	4a	4b
χ_1	1	1	1	1	1	1
χ_2	1	1	1	1	-1	-1
χ_3	2	2	-1	-1	0	0
χ_4	1	-1	1	-1	$E(4)$	$-E(4)$
χ_5	1	-1	1	-1	$-E(4)$	$E(4)$
χ_6	2	-2	-1	1	0	0

Trivial source character table of $G \cong \text{C3} : \text{C4}$ at $p = 3$:

Normalisers N_i	N_1				N_2			
p -subgroups of G up to conjugacy in G	P_1				P_2			
Representatives $n_j \in N_i$	1a	4a	2a	4b	1a	4a	2a	4b
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6$	3	1	3	1	0	0	0	0
$0 \cdot \chi_1 + 1 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6$	3	-1	3	-1	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 1 \cdot \chi_6$	3	$E(4)$	-3	$-E(4)$	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6$	3	$-E(4)$	-3	$E(4)$	0	0	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6$	1	1	1	1	1	1	1	1
$0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6$	1	-1	1	-1	1	-1	1	-1
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6$	1	$E(4)$	-1	$-E(4)$	1	$E(4)$	-1	$-E(4)$
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 0 \cdot \chi_6$	1	$-E(4)$	-1	$E(4)$	1	$-E(4)$	-1	$E(4)$

$$P_1 = \text{Group}([\langle \rangle]) \cong 1$$

$$P_2 = \text{Group}([(1, 8, 4)(2, 10, 6)(3, 11, 7)(5, 12, 9)]) \cong \text{C3}$$

$$N_1 = \text{Group}([(1, 2, 3, 5)(4, 10, 7, 12)(6, 11, 9, 8), (1, 3)(2, 5)(4, 7)(6, 9)(8, 11)(10, 12), (1, 4, 8)(2, 6, 10)(3, 7, 11)(5, 9, 12)]) \cong \text{C3} : \text{C4}$$

$$N_2 = \text{Group}([(1, 8, 4)(2, 10, 6)(3, 11, 7)(5, 12, 9), (1, 2, 3, 5)(4, 10, 7, 12)(6, 11, 9, 8)]) \cong \text{C3} : \text{C4}$$