

The group G is isomorphic to the group labelled by [12, 2] in the Small Groups library.

Ordinary character table of $G \cong \text{C12}$:

	1a	4a	2a	4b	3a	12a	6a	12b	3b	12c	6b	12d
χ_1	1	1	1	1	1	1	1	1	1	1	1	1
χ_2	1	-1	1	-1	1	-1	1	-1	1	-1	1	-1
χ_3	1	1	1	1	$E(3)$	$E(3)$	$E(3)$	$E(3)$	$E(3)^2$	$E(3)^2$	$E(3)^2$	$E(3)^2$
χ_4	1	-1	1	-1	$E(3)$	$-E(3)$	$E(3)$	$-E(3)$	$E(3)^2$	$-E(3)^2$	$E(3)^2$	$-E(3)^2$
χ_5	1	1	1	1	$E(3)^2$	$E(3)^2$	$E(3)^2$	$E(3)^2$	$E(3)$	$E(3)$	$E(3)$	$E(3)$
χ_6	1	-1	1	-1	$E(3)^2$	$-E(3)^2$	$E(3)^2$	$-E(3)^2$	$E(3)$	$-E(3)$	$E(3)$	$-E(3)$
χ_7	1	$E(4)$	-1	$-E(4)$	1	$E(4)$	-1	$-E(4)$	1	$E(4)$	-1	$-E(4)$
χ_8	1	$-E(4)$	-1	$E(4)$	1	$-E(4)$	-1	$E(4)$	1	$-E(4)$	-1	$E(4)$
χ_9	1	$E(4)$	-1	$-E(4)$	$E(3)$	$E(12)^7$	$-E(3)$	$-E(12)^7$	$E(3)^2$	$E(12)^{11}$	$-E(3)^2$	$-E(12)^{11}$
χ_{10}	1	$-E(4)$	-1	$E(4)$	$E(3)$	$-E(12)^7$	$-E(3)$	$E(12)^7$	$E(3)^2$	$-E(12)^{11}$	$-E(3)^2$	$E(12)^{11}$
χ_{11}	1	$E(4)$	-1	$-E(4)$	$E(3)^2$	$E(12)^{11}$	$-E(3)^2$	$-E(12)^{11}$	$E(3)$	$E(12)^7$	$-E(3)$	$-E(12)^7$
χ_{12}	1	$-E(4)$	-1	$E(4)$	$E(3)^2$	$-E(12)^{11}$	$-E(3)^2$	$E(12)^{11}$	$E(3)$	$-E(12)^7$	$-E(3)$	$E(12)^7$

Trivial source character table of $G \cong \text{C12}$ at $p = 2$:

Normalisers N_i	N_1			N_2			N_3		
p -subgroups of G up to conjugacy in G	P_1			P_2			P_3		
Representatives $n_j \in N_i$	1a	3a	3b	1a	3a	3b	1a	3a	3b
$1 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 1 \cdot \chi_7 + 1 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	4	4	4	0	0	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 1 \cdot \chi_9 + 1 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	4	$4 * E(3)$	$4 * E(3)^2$	0	0	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 1 \cdot \chi_{11} + 1 \cdot \chi_{12}$	4	$4 * E(3)^2$	$4 * E(3)$	0	0	0	0	0	0
$1 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	2	2	2	2	2	2	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	2	$2 * E(3)^2$	$2 * E(3)$	2	$2 * E(3)^2$	$2 * E(3)$	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	2	$2 * E(3)$	$2 * E(3)^2$	2	$2 * E(3)$	$2 * E(3)^2$	0	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	1	1	1	1	1	1	1	1	1
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	1	$E(3)$	$E(3)^2$	1	$E(3)$	$E(3)^2$	1	$E(3)$	$E(3)^2$
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12}$	1	$E(3)^2$	$E(3)$	1	$E(3)^2$	$E(3)$	1	$E(3)^2$	$E(3)$

$$P_1 = \text{Group}([()]) \cong 1$$

$$P_2 = \text{Group}([(4, 6)(5, 7)]) \cong \text{C2}$$

$$P_3 = \text{Group}([(4, 5, 6, 7), (4, 6)(5, 7)]) \cong \text{C4}$$

$$N_1 = \text{Group}([(1, 2, 3), (4, 5, 6, 7)]) \cong \text{C12}$$

$$N_2 = \text{Group}([(1, 2, 3), (4, 5, 6, 7)]) \cong \text{C12}$$

$$N_3 = \text{Group}([(1, 2, 3), (4, 5, 6, 7)]) \cong \text{C12}$$