

The group G is isomorphic to the group labelled by [40, 5] in the Small Groups library.
 Ordinary character table of $G \cong C_4 \times D_{10}$:

	1a	2a	4a	2b	5a	4b	2c	4c	20a	10a	5b	4d	20b	20c	10b	20d
χ_1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
χ_2	1	-1	-1	1	1	1	-1	-1	-1	1	1	-1	-1	1	-1	-1
χ_3	1	-1	1	1	1	-1	-1	1	1	1	-1	1	1	1	1	1
χ_4	1	1	-1	1	1	-1	1	-1	-1	1	1	-1	-1	1	-1	-1
χ_5	1	-1	$-E(4)$	-1	1	$E(4)$	1	$E(4)$	$-E(4)$	-1	1	$-E(4)$	$E(4)$	$-E(4)$	-1	$E(4)$
χ_6	1	-1	$E(4)$	-1	1	$-E(4)$	1	$-E(4)$	$E(4)$	-1	1	$E(4)$	$-E(4)$	$E(4)$	-1	$-E(4)$
χ_7	1	1	$-E(4)$	-1	1	$-E(4)$	-1	$E(4)$	$-E(4)$	-1	1	$E(4)$	$E(4)$	$-E(4)$	-1	$E(4)$
χ_8	1	1	$E(4)$	-1	1	$E(4)$	-1	$-E(4)$	$E(4)$	-1	1	$-E(4)$	$E(4)$	$-E(4)$	-1	$-E(4)$
χ_9	2	0	-2	2	$E(5)^2 + E(5)^3$	0	0	-2	$-E(5)^2 - E(5)^3$	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	$-E(5)^2 - E(5)^3$	$-E(5) - E(5)^4$	$E(5) + E(5)^4$	$-E(5) - E(5)^4$
χ_{10}	2	0	-2	2	$E(5) + E(5)^4$	0	0	-2	$-E(5) - E(5)^4$	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	$-E(5) - E(5)^4$	$-E(5)^2 - E(5)^3$	$E(5)^2 + E(5)^3$	$-E(5)^2 - E(5)^3$
χ_{11}	2	0	2	2	$E(5)^2 + E(5)^3$	0	0	2	$E(5)^2 + E(5)^3$	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	$E(5) + E(5)^4$	$E(5) + E(5)^4$
χ_{12}	2	0	2	2	$E(5) + E(5)^4$	0	0	2	$E(5) + E(5)^4$	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	$E(5)^2 + E(5)^3$	$E(5)^2 + E(5)^3$
χ_{13}	2	0	$-2 * E(4)$	-2	$E(5)^2 + E(5)^3$	0	0	$2 * E(4)$	$-E(20)^{13} - E(20)^{17}$	$-E(5)^2 - E(5)^3$	$E(5) + E(5)^4$	0	$E(20)^{13} + E(20)^{17}$	$-E(20) - E(20)^9$	$-E(5) - E(5)^4$	$E(20) + E(20)^9$
χ_{14}	2	0	$-2 * E(4)$	-2	$E(5) + E(5)^4$	0	0	$2 * E(4)$	$-E(20) - E(20)^9$	$-E(5) - E(5)^4$	$E(5)^2 + E(5)^3$	0	$E(20) + E(20)^9$	$-E(20)^{13} - E(20)^{17}$	$-E(5)^2 - E(5)^3$	$E(20)^{13} + E(20)^{17}$
χ_{15}	2	0	$2 * E(4)$	-2	$E(5)^2 + E(5)^3$	0	0	$-2 * E(4)$	$E(20)^{13} + E(20)^{17}$	$-E(5)^2 - E(5)^3$	$E(5) + E(5)^4$	0	$-E(20)^{13} - E(20)^{17}$	$E(20) + E(20)^9$	$-E(5) - E(5)^4$	$-E(20) - E(20)^9$
χ_{16}	2	0	$2 * E(4)$	-2	$E(5) + E(5)^4$	0	0	$-2 * E(4)$	$E(20) + E(20)^9$	$-E(5) - E(5)^4$	$E(5)^2 + E(5)^3$	0	$-E(20) - E(20)^9$	$E(20)^{13} + E(20)^{17}$	$-E(5)^2 - E(5)^3$	$-E(20)^{13} - E(20)^{17}$

Trivial source character table of $G \cong C_4 \times D_{10}$ at $p = 2$:

Normalisers N_i	N_1	N_2	N_3	N_4	N_5	N_6	N_7	N_8
p -subgroups of G up to conjugacy in G	P_1	P_2	P_3	P_4	P_5	P_6	P_7	P_8
Representatives $n_j \in N_i$								
1 · $\chi_1 + \chi_2 + 1 \cdot \chi_3 + 1 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7 + 1 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	1a	5a	5b	1a	1a	1a	1a	1a
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 1 \cdot \chi_9 + 0 \cdot \chi_{10} + 1 \cdot \chi_{11} + 0 \cdot \chi_{12} + 1 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	8	8	8	0	0	0	0	0
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 1 \cdot \chi_{10} + 0 \cdot \chi_{11} + 1 \cdot \chi_{12} + 0 \cdot \chi_{13} + 1 \cdot \chi_{14} + 0 \cdot \chi_{15} + 1 \cdot \chi_{16}$	8	$4 * E(5) + 4 * E(5)^4$	$4 * E(5) + 4 * E(5)^4$	0	0	0	0	0
1 · $\chi_1 + 1 \cdot \chi_2 + 1 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	4	4	4	4	4	0	0	0
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 1 \cdot \chi_9 + 0 \cdot \chi_{10} + 1 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	4	$2 * E(5)^2 + 2 * E(5)^3$	$2 * E(5) + 2 * E(5)^4$	4	$2 * E(5)^2 + 2 * E(5)^3$	$2 * E(5) + 2 * E(5)^4$	0	0
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 1 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	4	$2 * E(5) + 2 * E(5)^4$	$2 * E(5)^2 + 2 * E(5)^3$	4	$2 * E(5) + 2 * E(5)^4$	$2 * E(5)^2 + 2 * E(5)^3$	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 1 \cdot \chi_7 + 1 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	4	4	4	0	0	0	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 1 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	4	4	4	0	0	0	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	2	2	2	2	2	0	2	2
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 1 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	2	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	2	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	0
0 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 1 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	2	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	2	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	2	2	2	2	2	0	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	2	2	2	2	2	0	0	0
1 · $\chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} + 0 \cdot \chi_{14} + 0 \cdot \chi_{15} + 0 \cdot \chi_{16}$	1	1	1	1	1	1	1	1

$P_1 = \text{Group}([()]) \cong 1$

$P_2 = \text{Group}([(1, 4)(2, 7)(3, 9)(5, 11)(6, 13)(8, 15)(10, 17)(12, 19)(14, 21)(16, 23)(18, 25)(20, 27)(22, 29)(24, 31)(26, 33)(28, 35)(30, 36)(32, 38)(34, 39)(37, 40)]) \cong C_2$