

The group G is isomorphic to the group labelled by [52, 3] in the Small Groups library.

Ordinary character table of $G \cong \text{C13} : \text{C4}$:

	$1a$	$13a$	$13b$	$13c$	$4a$	$2a$	$4b$
χ_1	1	1	1	1	1	1	1
χ_2	1	1	1	1	$E(4)$	-1	$-E(4)$
χ_3	1	1	1	1	-1	1	-1
χ_4	1	1	1	1	$-E(4)$	-1	$E(4)$
χ_5	4	$E(13) + E(13)^5 + E(13)^8 + E(13)^{12}$	$E(13)^2 + E(13)^3 + E(13)^{10} + E(13)^{11}$	$E(13)^4 + E(13)^6 + E(13)^7 + E(13)^9$	0	0	0
χ_6	4	$E(13)^2 + E(13)^3 + E(13)^{10} + E(13)^{11}$	$E(13)^4 + E(13)^6 + E(13)^7 + E(13)^9$	$E(13) + E(13)^5 + E(13)^8 + E(13)^{12}$	0	0	0
χ_7	4	$E(13)^4 + E(13)^6 + E(13)^7 + E(13)^9$	$E(13) + E(13)^5 + E(13)^8 + E(13)^{12}$	$E(13)^2 + E(13)^3 + E(13)^{10} + E(13)^{11}$	0	0	0

Trivial source character table of $G \cong \text{C13} : \text{C4}$ at $p = 13$:

Normalisers N_i	N_1				N_2			
p -subgroups of G up to conjugacy in G	P_1				P_2			
Representatives $n_j \in N_i$	$1a$	$4a$	$2a$	$4b$	$1a$	$4a$	$2a$	$4b$
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7$	13	-1	1	-1	0	0	0	0
$0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7$	13	$E(4)$	-1	$-E(4)$	0	0	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7$	13	1	1	1	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 1 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7$	13	$-E(4)$	-1	$E(4)$	0	0	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7$	1	1	1	1	1	1	1	1
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7$	1	-1	1	-1	1	-1	1	-1
$0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7$	1	$E(4)$	-1	$-E(4)$	1	$E(4)$	-1	$-E(4)$
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7$	1	$-E(4)$	-1	$E(4)$	1	$-E(4)$	-1	$E(4)$

$$P_1 = \text{Group}([\langle \rangle]) \cong 1$$

$$P_2 = \text{Group}([(1, 36, 20, 4, 40, 24, 8, 44, 28, 12, 48, 32, 16)(2, 38, 22, 6, 42, 26, 10, 46, 30, 14, 50, 34, 18)(3, 39, 23, 7, 43, 27, 11, 47, 31, 15, 51, 35, 19)(5, 41, 25, 9, 45, 29, 13, 49, 33, 17, 52, 37, 21)]) \cong \text{C13}$$

$$N_1 = \text{Group}([(1, 2, 3, 5)(4, 34, 51, 25)(6, 35, 52, 20)(7, 37, 48, 22)(8, 14, 47, 45)(9, 32, 50, 23)(10, 15, 49, 40)(11, 17, 44, 42)(12, 46, 43, 13)(16, 26, 39, 33)(18, 27, 41, 28)(19, 29, 36, 30)(21, 24, 38, 31), (1, 3)(2, 5)(4, 51)(6, 52)(7, 48)(8, 47)(9, 50)(10, 49)(11, 44)(12, 43)(13, 46)(14, 45)(15, 40)(16, 39)(17, 42)(18, 41)(19, 36)(20, 35)(21, 38)(22, 37)(23, 32)(24, 31)(25, 34)(26, 33)(27, 28)(29, 30), (1, 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48)(2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50)(3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 51)(5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 52)]) \cong \text{C13} : \text{C4}$$

$$N_2 = \text{Group}([(1, 36, 20, 4, 40, 24, 8, 44, 28, 12, 48, 32, 16)(2, 38, 22, 6, 42, 26, 10, 46, 30, 14, 50, 34, 18)(3, 39, 23, 7, 43, 27, 11, 47, 31, 15, 51, 35, 19)(5, 41, 25, 9, 45, 29, 13, 49, 33, 17, 52, 37, 21), (1, 2, 3, 5)(4, 34, 51, 25)(6, 35, 52, 20)(7, 37, 48, 22)(8, 14, 47, 45)(9, 32, 50, 23)(10, 15, 49, 40)(11, 17, 44, 42)(12, 46, 43, 13)(16, 26, 39, 33)(18, 27, 41, 28)(19, 29, 36, 30)(21, 24, 38, 31)]) \cong \text{C13} : \text{C4}$$