

The group G is isomorphic to the group labelled by [660, 13] in the Small Groups library.

Ordinary character table of $G \cong \text{PSL}(2,11)$:

	1a	2a	3a	5a	5b	6a	11a	11b
χ_1	1	1	1	1	1	1	1	1
χ_2	5	1	-1	0	0	1	$E(11) + E(11)^3 + E(11)^4 + E(11)^5 + E(11)^9$	$E(11)^2 + E(11)^6 + E(11)^7 + E(11)^8 + E(11)^{10}$
χ_3	5	1	-1	0	0	1	$E(11)^2 + E(11)^6 + E(11)^7 + E(11)^8 + E(11)^{10}$	$E(11) + E(11)^3 + E(11)^4 + E(11)^5 + E(11)^9$
χ_4	10	-2	1	0	0	1	-1	-1
χ_5	10	2	1	0	0	-1	-1	-1
χ_6	11	-1	-1	1	1	-1	0	0
χ_7	12	0	0	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	1	1
χ_8	12	0	0	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	1	1

Trivial source character table of $G \cong \text{PSL}(2,11)$ at $p = 5$:

Normalisers N_i	N_1						N_2	
p -subgroups of G up to conjugacy in G	P_1						P_2	
Representatives $n_j \in N_i$	1a	2a	3a	6a	11a	11b	1a	2a
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 1 \cdot \chi_7 + 1 \cdot \chi_8$	25	1	1	1	3	3	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 1 \cdot \chi_6 + 1 \cdot \chi_7 + 1 \cdot \chi_8$	35	-1	-1	-1	2	2	0	0
$0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	5	1	-1	1	$E(11) + E(11)^3 + E(11)^4 + E(11)^5 + E(11)^9$	$E(11)^2 + E(11)^6 + E(11)^7 + E(11)^8 + E(11)^{10}$	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	5	1	-1	1	$E(11)^2 + E(11)^6 + E(11)^7 + E(11)^8 + E(11)^{10}$	$E(11) + E(11)^3 + E(11)^4 + E(11)^5 + E(11)^9$	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	10	-2	1	1	-1	-1	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	10	2	1	-1	-1	-1	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	1	1	1	1	1	1	1	1
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 1 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8$	11	-1	-1	-1	0	0	1	-1

$$P_1 = \text{Group}([()]) \cong 1$$

$$P_2 = \text{Group}([(2, 6, 8, 7, 10)(3, 5, 9, 4, 11)]) \cong \text{C5}$$

$$N_1 = \text{Group}([(2, 10)(3, 4)(5, 9)(6, 7), (1, 2, 11)(3, 5, 10)(6, 8, 9)]) \cong \text{PSL}(2,11)$$

$$N_2 = \text{Group}([(2, 6, 8, 7, 10)(3, 5, 9, 4, 11), (3, 11)(4, 5)(6, 10)(7, 8)]) \cong \text{D10}$$