

**Corollary 15.1**

Character values are algebraic integers.

**Proof:** By the above, roots of unity are algebraic integers. Since the algebraic integers form a ring, so are sums of roots of unity. Hence the claim follows from Property 7.4(b). ■

## 16 Central Characters

We now extend representations/characters of finite groups to "representations/characters" of the centre of the group algebra  $\mathbb{C}G$  in order to obtain further results on character values, which we will use in the next sections in order to prove Burnside's  $p^a q^b$  theorem.

**Definition 16.1 (Class sums)**

The elements  $\hat{C}_j := \sum_{g \in C_j} g \in \mathbb{C}G$  ( $1 \leq j \leq r$ ) are called the **class sums** of  $G$ .

**Lemma 16.2**

The class sums  $\{\hat{C}_j \mid 1 \leq j \leq r\}$  form a  $\mathbb{C}$ -basis of  $Z(\mathbb{C}G)$ . In other words,  $Z(\mathbb{C}G) = \bigoplus_{j=1}^r \mathbb{C}\hat{C}_j$ .

**Proof:** Notice that the class sums are clearly  $\mathbb{C}$ -linearly independent because the group elements are.

' $\supseteq$ ':  $\forall 1 \leq j \leq r$  and  $\forall g \in G$ , we have

$$g \cdot \hat{C}_j = g(g^{-1}\hat{C}_jg) = \hat{C}_j \cdot g.$$

Extending by  $\mathbb{C}$ -linearity, we get  $a \cdot \hat{C}_j = \hat{C}_j \cdot a \quad \forall 1 \leq j \leq r$  and  $\forall a \in \mathbb{C}G$ . Thus  $\bigoplus_{j=1}^r \mathbb{C}\hat{C}_j \subseteq Z(\mathbb{C}G)$ .

' $\subseteq$ ': Let  $a \in Z(\mathbb{C}G)$  and write  $a = \sum_{g \in G} \lambda_g g$  with  $\{\lambda_g\}_{g \in G} \in \mathbb{C}$ . Since  $a$  is central, for every  $h \in G$ , we have

$$\sum_{g \in G} \lambda_g g = a = hah^{-1} = \sum_{g \in G} \lambda_g (hgh^{-1}).$$

Comparing coefficients yield  $\lambda_g = \lambda_{hgh^{-1}} \quad \forall g, h \in G$ . Namely, the coefficients  $\lambda_g$  are constant on the conjugacy classes of  $G$ , and hence

$$a = \sum_{j=1}^r \lambda_{g_j} \hat{C}_j \in \bigoplus_{j=1}^r \mathbb{C}\hat{C}_j. \quad \blacksquare$$

Now, notice that by definition the class sums  $\hat{C}_j$  ( $1 \leq j \leq r$ ) are elements of the subring  $\mathbb{Z}G$  of  $\mathbb{C}G$ , hence of the centre of  $\mathbb{Z}G$ .

**Corollary 16.3**

- (a)  $Z(\mathbb{Z}G)$  is finitely generated as a  $\mathbb{Z}$ -module.
- (b) The centre  $Z(\mathbb{Z}G)$  of the group ring  $\mathbb{Z}G$  is integral over  $\mathbb{Z}$ ; in particular the class sums  $\hat{C}_j$  ( $1 \leq j \leq r$ ) are algebraic integers.

**Proof:**

- (a) It follows directly from the second part of the proof of Lemma 16.2 that the class sums  $\hat{C}_j$  ( $1 \leq j \leq r$ ) span  $Z(\mathbb{Z}G)$  as a  $\mathbb{Z}$ -module.
- (b) The centre  $Z(\mathbb{Z}G)$  is integral over  $\mathbb{Z}$  by Theorem D.2 because it is finitely generated as a  $\mathbb{Z}$ -module by (a). ■

**Notation 16.4 (Central characters)**

If  $\chi \in \text{Irr}(G)$ , then we may consider a  $\mathbb{C}$ -representation affording  $\chi$ , say  $\rho^\chi : G \rightarrow \text{GL}(\mathbb{C}^{n(\chi)}) = \text{Aut}_{\mathbb{C}}(\mathbb{C}^{n(\chi)})$  with  $n(\chi) := \chi(1)$ . This group homomorphism extends by  $\mathbb{C}$ -linearity to a  $\mathbb{C}$ -algebra homomorphism

$$\begin{aligned} \tilde{\rho}^\chi : \quad \mathbb{C}G &\longrightarrow \text{End}_{\mathbb{C}}(\mathbb{C}^{n(\chi)}) \\ a = \sum_{g \in G} \lambda_g g &\mapsto \tilde{\rho}^\chi(a) = \sum_{g \in G} \lambda_g \rho^\chi(g). \end{aligned}$$

Now, if  $z \in Z(\mathbb{C}G)$ , then for each  $g \in G$ , we have

$$\tilde{\rho}^\chi(z)\tilde{\rho}^\chi(g) = \tilde{\rho}^\chi(zg) = \tilde{\rho}^\chi(gz) = \tilde{\rho}^\chi(g)\tilde{\rho}^\chi(z).$$

As we have already seen in Chapter 2 on Schur's Lemma this means that  $\tilde{\rho}^\chi(z)$  is  $\mathbb{C}G$ -linear. This holds in particular if  $z$  is a class sum. Therefore, by Schur's Lemma, for each  $1 \leq j \leq r$  there exists a scalar  $\omega_\chi(\hat{C}_j) \in \mathbb{C}$  such that

$$\tilde{\rho}^\chi(\hat{C}_j) = \omega_\chi(\hat{C}_j) \cdot I_{n(\chi)}.$$

The functions defined by

$$\begin{aligned} \omega_\chi : \quad Z(\mathbb{C}G) &\longrightarrow \mathbb{C} \\ \hat{C}_j &\mapsto \omega_\chi(\hat{C}_j) \end{aligned}$$

and extended by  $\mathbb{C}$ -linearity to the whole of  $Z(\mathbb{C}G)$ , where  $\chi$  runs through  $\text{Irr}(G)$ , are called the **central characters** of  $\mathbb{C}G$  (or simply of  $G$ ).

**Remark 16.5**

If  $z \in Z(G)$ , then  $[z] = \{z\}$  and therefore the corresponding class sum is  $z$  itself. Therefore, we may see the functions  $\omega_\chi|_{Z(G)} : Z(G) \rightarrow \mathbb{C}$  as representations of  $Z(G)$  of degree 1, or equivalently as linear characters of  $Z(G)$ .

**Theorem 16.6 (Integrality Theorem)**

The values  $\omega_\chi(\hat{C}_j)$  ( $\chi \in \text{Irr}(G)$ ,  $1 \leq j \leq r$ ) of the central characters of  $G$  are algebraic integers. Moreover,

$$\omega_\chi(\hat{C}_j) = \frac{|C_j|}{\chi(1)} \chi(g_j) \quad \forall \chi \in \text{Irr}(G), \forall 1 \leq j \leq r.$$

**Proof:** Let  $\chi \in \text{Irr}(G)$  and  $1 \leq j \leq r$ . By Corollary 16.3 the class sum  $\hat{C}_j$  is an algebraic integer. Thus there exist integers  $n \in \mathbb{Z}_{>0}$  and  $a_0, \dots, a_{n-1} \in \mathbb{Z}$  such that  $\hat{C}_j^n + a_{n-1}\hat{C}_j^{n-1} + \dots + a_0 = 0$ . Applying  $\omega_\chi$  yields  $\omega_\chi(\hat{C}_j)^n + a_{n-1}\omega_\chi(\hat{C}_j)^{n-1} + \dots + a_0 = \omega_\chi(0) = 0$ , so that  $\omega_\chi(\hat{C}_j)$  is also an algebraic integer. Now, according to Notation 16.4 we have

$$\chi(1)\omega_\chi(\hat{C}_j) = \text{Tr}(\tilde{\rho}^\chi(\hat{C}_j)) = \text{Tr}\left(\sum_{g \in C_j} \rho^\chi(g)\right) = \sum_{g \in C_j} \text{Tr}(\rho^\chi(g)) = \sum_{g \in C_j} \chi(g) = |C_j|\chi(g),$$

where the last equality holds because characters are class functions. The claim follows. ■

**Corollary 16.7**

If  $\chi \in \text{Irr}(G)$ , then  $\chi(1)$  divides  $|G|$ .

**Proof:** By the 1st Orthogonality Relations we have

$$\frac{|G|}{\chi(1)} = \frac{|G|}{\chi(1)} \langle \chi, \chi \rangle_G = \frac{1}{\chi(1)} \sum_{g \in G} \chi(g)\chi(g^{-1}) = \frac{1}{\chi(1)} \sum_{j=1}^r |C_j| \chi(g_j)\chi(g_j^{-1}) = \sum_{j=1}^r \underbrace{\frac{|C_j|}{\chi(1)}}_{\omega_\chi(\hat{C}_j)} \chi(g_j)\chi(g_j^{-1}).$$

Now, for each  $1 \leq j \leq r$ ,  $\omega_\chi(g_j)$  is an algebraic integer by the Integrality Theorem and  $\chi(g_j^{-1})$  is an algebraic integer by Corollary 15.1. Hence  $|G|/\chi(1)$  is an algebraic integer because these form a subring of  $\mathbb{C}$ . Moreover, clearly  $|G|/\chi(1) \in \mathbb{Q}$ . As the algebraic integers in  $\mathbb{Q}$  are just the elements of  $\mathbb{Z}$ , we obtain that  $|G|/\chi(1) \in \mathbb{Z}$ , as claimed. ■

**Example 8 (The degrees of the irreducible characters of  $\text{GL}_3(\mathbb{F}_2)$ )**

The group  $G := \text{GL}_3(\mathbb{F}_2)$  is a simple group of order

$$|G| = \# \mathbb{F}_2\text{-bases of } \mathbb{F}_2^3 = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 168 = 2^3 \cdot 3 \cdot 7.$$

For the purpose of this example we accept without proof that  $G$  is simple and that it has 6 conjugacy classes.

**Question:** can we compute the degrees of the irreducible characters of  $\text{GL}_3(\mathbb{F}_2)$ ?

(1) By the above  $|\text{Irr}(G)| = |C(G)| = 6$  and the degree formula yields:

$$1 + \sum_{i=2}^6 \chi_i(1)^2 = |G| = 168.$$

(2) Next, as  $G$  is simple non-abelian,  $G = G'$  and therefore  $G$  has  $|G : G'| = 1$  linear characters by Corollary 14.8, namely

$$\chi_i(1) \geq 2 \text{ for each } 2 \leq i \leq 6.$$

Thus, at this stage, we would have the following possibilities for the degrees of the 6 irreducible characters of  $G$ :

$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	9
1	2	3	3	8	9
1	2	5	5	7	8
1	2	4	7	7	7
1	3	3	6	7	8

(3) By Corollary 16.7 we now know that  $\chi_i(1) \mid |G|$  for each  $2 \leq i \leq 6$ . Therefore, as  $5 \nmid |G|$  and  $9 \nmid |G|$ , the first three rows can already be discarded:

$\chi_1(1)$	$\chi_2(1)$	$\chi_3(1)$	$\chi_4(1)$	$\chi_5(1)$	$\chi_6(1)$
1	2	4	5	6	<del>9</del>
1	2	3	3	8	<del>9</del>
1	2	<del>5</del>	<del>5</del>	7	8
1	2	4	7	7	7
1	3	3	6	7	8

(4) In order to eliminate the last-but-one possibility, we apply [Exercise 21(b), Sheet 6] saying that a simple group cannot have an irreducible character of degree 2. Hence

$$\chi_1(1) = 1, \chi_2(1) = 3, \chi_3(1) = 3, \chi_4(1) = 6, \chi_5(1) = 7, \chi_6(1) = 8.$$

## 17 The Centre of a Character

### Definition 17.1 (Centre of a character)

The centre of a character  $\chi$  of  $G$  is  $Z(\chi) := \{g \in G \mid |\chi(g)| = \chi(1)\}$ .

**Note:** Recall that in contrast,  $\chi(g) = \chi(1) \iff g \in \ker(\chi)$ .

### Example 9

Recall from Example 5 that the character table of  $G = S_3$  is

	Id	(12)	(123)
$\chi_1$	1	1	1
$\chi_2$	1	-1	1
$\chi_3$	2	0	-1

Hence  $Z(\chi_1) = Z(\chi_2) = G$  and  $Z(\chi_3) = \{\text{Id}\}$ .

### Lemma 17.2

If  $\rho : G \rightarrow \text{GL}(V)$  is a  $\mathbb{C}$ -representation with character  $\chi$  and  $g \in G$ , then:

$$|\chi(g)| = \chi(1) \iff \rho(g) \in \mathbb{C}^\times \text{Id}_V.$$

In other words  $Z(\chi) = \rho^{-1}(\mathbb{C}^\times \text{Id}_V)$ .

**Proof:** Let  $n := \chi(1)$ . Recall that we can find a  $\mathbb{C}$ -basis  $B$  of  $V$  such that  $(\rho(g))_B$  is a diagonal matrix with diagonal entries  $\varepsilon_1, \dots, \varepsilon_n$  which are  $\sigma(g)$ -th roots of unity. Hence  $\varepsilon_1, \dots, \varepsilon_n$  are the eigenvalues of  $\rho(g)$ . Applying the Cauchy-Schwartz inequality to the vectors  $v := (\varepsilon_1, \dots, \varepsilon_n)$  and  $w := (1, \dots, 1)$  in  $\mathbb{C}^n$  yields

$$|\chi(g)| = |\varepsilon_1 + \dots + \varepsilon_n| = |\langle v, w \rangle| \leq \|v\| \|w\| = \sqrt{n} \sqrt{n} = n = \chi(1)$$

and equality implies that  $v$  and  $w$  are  $\mathbb{C}$ -linearly dependent so that  $\varepsilon_1 = \dots = \varepsilon_n =: \varepsilon$ . Therefore  $\rho(g) \in \mathbb{C}^\times \text{Id}_V$ . Conversely, if  $\rho(g) \in \mathbb{C}^\times \text{Id}_V$ , then there exists  $\lambda \in \mathbb{C}^\times$  such that  $\rho(g) = \lambda \text{Id}_V$ . Therefore the eigenvalues of  $\rho(g)$  are all equal to  $\lambda$ , i.e.  $\lambda = \varepsilon_1 = \dots = \varepsilon_n$  and therefore

$$|\chi(g)| = |n\lambda| = n|\lambda| = n \cdot 1 = n.$$

