
Chapter 6. The Mackey Formula and Clifford Theory

The results in this chapter go more deeply into the theory. We start with the so-called *Mackey decomposition formula*, which provides us with yet another relationship between induction and restriction. After that we explain Clifford's theorem, which considers restriction/induction of simple modules to/from a normal subgroup. These results are essential and have many consequences throughout representation theory of finite groups.

Notation. Throughout this chapter, unless otherwise specified, we let G denote a finite group and let (F, \mathcal{O}, k) be a p -modular system, which is splitting for G and all its subgroups. We let $K \in \{F, \mathcal{O}, k\}$ and assume all KG -modules considered are assumed are **free of finite rank as K -modules** (hence **finitely generated** as KG -modules).

References:

- [Alp86] J. L. Alperin. *Local representation theory*. Vol. 11. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1986.
- [Ben98] D. J. Benson. *Representations and cohomology. I*. Vol. 30. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1998.
- [CR90] C. W. Curtis and I. Reiner. *Methods of representation theory. Vol. I*. John Wiley & Sons, Inc., New York, 1990.
- [LP10] K. Lux and H. Pahlings. *Representations of groups*. Vol. 124. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2010.
- [Web16] P. Webb. *A course in finite group representation theory*. Vol. 161. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2016.

18 Double cosets

Definition 18.1 (*Double cosets*)

Given subgroups H and L of G we define for each $g \in G$

$$HgL := \{hg\ell \in G \mid h \in H, \ell \in L\}$$

and call this subset of G the (H, L) -**double coset** of g . Moreover, we let $H \backslash G / L$ denote the set of all (H, L) -double cosets of G .

First, we want to prove that the (H, L) -double cosets partition the group G .

Lemma 18.2

Let $H, L \leq G$.

- (a) Each (H, L) -double coset is a disjoint union of right cosets of H and a disjoint union of left cosets of L .
- (b) Any two (H, L) -double cosets either coincide or are disjoint. Hence, letting $[H \backslash G / L]$ denote a set of representatives for the (H, L) -double cosets of G , we have

$$G = \bigsqcup_{g \in [H \backslash G / L]} HgL.$$

Proof:

- (a) If $hgl \in HgL$ and $\ell_1 \in L$, then $hgl \cdot \ell_1 = hg(\ell\ell_1) \in HgL$. It follows that the entire left coset of L that contains hgl is contained in HgL . This proves that HgL is a union of left cosets of L . A similar argument proves that HgL is a union of right cosets of H .
- (b) Let $g_1, g_2 \in G$. If $h_1g_1\ell_1 = h_2g_2\ell_2 \in Hg_1L \cap Hg_2L$, then $g_1 = h_1^{-1}h_2g_2\ell_2\ell_1^{-1} \in Hg_2L$ so that $Hg_1L \subseteq Hg_2L$. Similarly $Hg_2L \subseteq Hg_1L$. Thus if two double cosets are not disjoint, they coincide. ■

If X is a left G -set we use the standard notation $G \backslash X$ for the set of orbits of G on X , and denote a set of representatives for these orbits by $[G \backslash X]$. Similarly if Y is a right G -set we write Y/G and $[Y/G]$. We shall also repeatedly use the orbit-stabiliser theorem without further mention: in other words, if X is a transitive left G -set and $x \in X$ then $X \cong G/\text{Stab}_G(x)$ (i.e. the set of left cosets of the stabiliser of x in G), and similarly for right G -sets.

Exercise 18.3

- (a) Let $H, L \leq G$. Prove that the set of (H, L) -double cosets is in bijection with the set of orbits $H \backslash (G/L)$, and also with the set of orbits $(H \backslash G)/L$ under the mappings

$$HgL \mapsto H(gL) \in H \backslash (G/L)$$

$$HgL \mapsto (Hg)L \in (H \backslash G)/L.$$

This justifies the notation $H \backslash G / L$ for the set of (H, L) -double cosets.

- (b) Let $G = S_3$. Consider $H = L := S_2 = \{\text{Id}, (1\ 2)\}$ as a subgroup of S_3 . Prove that

$$[S_2 \backslash S_3 / S_2] = \{\text{Id}, (1\ 2\ 3)\}$$

while

$$S_2 \backslash S_3 / S_2 = \{\{\text{Id}, (1\ 2)\}, \{(1\ 2\ 3), (1\ 3\ 2), (1\ 3), (2\ 3)\}\}.$$

19 The Mackey formula

If H and L are subgroups of G , we wish to describe what happens if we induce a KL -module from L to G and then restrict it to H .

Remark 19.1

We need to examine KG regarded as a (KH, KL) -bimodule (i.e. with left and right external laws by multiplication in G). Since $G = \bigsqcup_{g \in [H \backslash G/L]} HgL$, we have

$$KG = \bigoplus_{g \in [H \backslash G/L]} K \langle HgL \rangle$$

as (KH, KL) -bimodule, where $K \langle HgL \rangle$ denotes the free K -module with K -basis HgL .

Now if M is a KL -module, we will also write ${}^g M$ for $g \otimes M$, which is a left $K({}^g L)$ -module with

$$(g\ell g^{-1}) \cdot (g \otimes m) = g \otimes \ell m$$

for each $\ell \in L$ and each $m \in M$. With this notation, we have

$$K \langle HgL \rangle \cong KH \otimes_{K(H \cap {}^g L)} (g \otimes KL),$$

where $hg\ell \in HgL$ corresponds to $h \otimes g \otimes \ell$.

Theorem 19.2 (Mackey formula)

Let $H, L \leq G$ and let M be a KL -module. Then, as KH -modules,

$$M \uparrow_L^G \downarrow_H^G \cong \bigoplus_{g \in [H \backslash G/L]} ({}^g M \downarrow_{H \cap {}^g L}^{{}^g L}) \uparrow_{H \cap {}^g L}^H \cdot$$

Proof: It follows from Remark 19.1 that as left KH -modules we have

$$\begin{aligned} M \uparrow_L^G \downarrow_H^G &\cong (KG \otimes_{KL} M) \downarrow_H^G \cong \bigoplus_{g \in [H \backslash G/L]} K \langle HgL \rangle \otimes_{KL} M \\ &\cong \bigoplus_{g \in [H \backslash G/L]} KH \otimes_{K(H \cap {}^g L)} (g \otimes KL) \otimes_{KL} M \\ &\cong \bigoplus_{g \in [H \backslash G/L]} KH \otimes_{K(H \cap {}^g L)} (g \otimes M) \downarrow_{H \cap {}^g L}^{{}^g L} \\ &\cong \bigoplus_{g \in [H \backslash G/L]} ({}^g M \downarrow_{H \cap {}^g L}^{{}^g L}) \uparrow_{H \cap {}^g L}^H \cdot \end{aligned}$$

■

Remark 19.3

Given an arbitrary finite group Z , write ${}_{KZ} \mathbf{mod}$ for the category of KZ -modules which are free of finite rank as K -modules. Then, expressed in categorical terms, the Mackey formula says that we have the following equality of functors from ${}_{KL} \mathbf{mod}$ to ${}_{KH} \mathbf{mod}$:

$$\text{Res}_H^G \circ \text{Ind}_L^G = \bigoplus_{g \in [H \backslash G/L]} \text{Ind}_{H \cap {}^g L}^H \circ \text{Res}_{H \cap {}^g L}^{{}^g L} \circ \text{Inn}(g)$$

where $\text{Inn}(g)$ is conjugation by $g \in G$.

Exercise 19.4

Let $H, L \leq G$, let M be a KL -module and let N be a KH -module. Use the Mackey formula to prove that:

$$(a) \quad M \uparrow_L^G \otimes_K N \uparrow_H^G \cong \bigoplus_{g \in [H \backslash G / L]} ({}^g M \downarrow_{H \cap {}^g L}^L \otimes_K N \downarrow_{H \cap {}^g L}^H) \uparrow_{H \cap {}^g L}^G;$$

$$(b) \quad \text{Hom}_K(M \uparrow_L^G, N \uparrow_H^G) \cong \bigoplus_{g \in [H \backslash G / L]} (\text{Hom}_K({}^g M \downarrow_{H \cap {}^g L}^L, N \downarrow_{H \cap {}^g L}^H)) \uparrow_{H \cap {}^g L}^G.$$

20 Clifford theory

We now turn to *Clifford's theorem*, which we present in a weak and a strong form. Clifford theory is a collection of results about induction and restriction of simple modules from/to normal subgroups.

Throughout this section, we assume that K is one of F or k .

Theorem 20.1 (Clifford's Theorem, weak form)

If $U \trianglelefteq G$ is a normal subgroup and S is a simple KG -module, then $S \downarrow_U^G$ is semisimple.

Proof: Let V be any simple KU -submodule of $S \downarrow_U^G$. Now, notice that for every $g \in G$, $gV := \{g v \mid v \in V\}$ is also a KU -submodule of $S \downarrow_U^G$, since $U \trianglelefteq G$ for any $u \in U$ we have

$$u \cdot gV = g \cdot \underbrace{(g^{-1} u g)}_{\in U} V = gV$$

Moreover, gV is also simple, since if W were a non-trivial proper KU -submodule of gV then $g^{-1}W$ would also be a non-trivial proper submodule of $g^{-1}gV = V$. Now $\sum_{g \in G} gV$ is non-zero and it is a KG -submodule of S , which is simple, hence $\sum_{g \in G} gV = S$. Restricting to U , we obtain that

$$S \downarrow_U^G = \sum_{g \in G} gV$$

is a sum of simple KU -submodules. Hence $S \downarrow_U^G$ is semisimple. ■

Remark 20.2

The KU -submodules gV which appear in the proof of Theorem 20.1 are isomorphic to modules we have seen before! More precisely the map

$$\begin{aligned} g \otimes V &\longrightarrow gV \\ g \otimes v &\longmapsto gv \end{aligned}$$

is a KU -isomorphism, since $U \trianglelefteq G$ implies that ${}^g U = U$ and hence the action of U on $g \otimes V$ (see Remark 19.1) and gV is prescribed in the same way.

Theorem 20.3 (Clifford's Theorem, strong form)

Let $U \trianglelefteq G$ be a normal subgroup and let S be a simple KG -module. Then we may write

$$S \downarrow_U^G = S_1^{a_1} \oplus \dots \oplus S_r^{a_r}$$

where $r \in \mathbb{Z}_{>0}$ and S_1, \dots, S_r are pairwise non-isomorphic simple KU -modules, occurring with multiplicities a_1, \dots, a_r respectively. Moreover, the following statements hold:

- (i) the group G permutes the homogeneous components of $S \downarrow_U^G$ transitively;
- (ii) $a_1 = a_2 = \dots = a_r$ and $\dim_K(S_1) = \dots = \dim_K(S_r)$; and
- (iii) $S \cong (S_1^{a_1}) \uparrow_{H_1}^G$ as KG -modules, where $H_1 = \text{Stab}_G(S_1^{a_1})$.

Proof: The fact that $S \downarrow_U^G$ is semisimple and hence can be written as a direct sum as claimed follows from Theorem 20.1. Moreover, by the chapter on semisimplicity of rings and modules, we know that for each $1 \leq i \leq r$ the homogeneous component $S_i^{a_i}$ is characterised by Proposition 7.1: it is the unique largest KU -submodule which is isomorphic to a direct sum of copies of S_i .

Now, if $g \in G$ then $g(S_i^{a_i}) = (gS_i)^{a_i}$, where gS_i is a simple KU -submodule of $S \downarrow_U^G$ (see the proof of the weak form of Clifford's Theorem). Hence there exists an index $1 \leq j \leq r$ such that $gS_i = S_j$ and $g(S_i^{a_i}) \subseteq S_j^{a_j}$ (alternatively to Proposition 7.1, the theorem of Krull-Schmidt can also be invoked here). Because $\dim_K(S_i) = \dim_K(gS_i)$, we have that $a_i \leq a_j$. Similarly, since $S_j = g^{-1}S_i$, we obtain $a_j \leq a_i$. Hence $a_i = a_j$ holds. Because

$$S \downarrow_U^G = g(S \downarrow_U^G) = g(S_1^{a_1}) \oplus \dots \oplus g(S_r^{a_r}),$$

we actually have that G permutes the homogeneous components. Moreover, as $\sum_{g \in G} g(S_1^{a_1})$ is a non-zero KG -submodule of S , which is simple, we have that $\sum_{g \in G} g(S_1^{a_1}) = S$, and so the action on the homogeneous components is transitive. This establishes both (i) and (ii).

For (iii), we define a K -homomorphism via the map

$$\begin{aligned} \Phi : (S_1^{a_1}) \uparrow_{H_1}^G = KG \otimes_{KH_1} S_1^{a_1} &= \bigoplus_{g \in [G/H_1]} g \otimes S_1^{a_1} &\longrightarrow S \\ g \otimes m & &\mapsto gm \end{aligned}$$

that is, where $g \otimes m \in g \otimes S_1^{a_1}$. This is in fact a KG -homomorphism. Furthermore, the K -subspaces $g(S_1^{a_1})$ of S are in bijection with the cosets G/H_1 , since G permutes them transitively by (i), and the stabiliser of one of them is H_1 . Thus both $KG \otimes_{KH_1} S_1^{a_1}$ and S are the direct sum of $|G : H_1|$ K -subspaces $g \otimes S_1^{a_1}$ and $g(S_1^{a_1})$ respectively, each K -isomorphic to $S_1^{a_1}$ (via $g \otimes m \leftrightarrow m$ and $gm \leftrightarrow m$). Thus the restriction of Φ to each summand is an isomorphism, and so Φ itself must be bijective, hence a KG -isomorphism. ■

One application of Clifford's theory is for example the following Corollary:

Corollary 20.4

If ℓ is a prime number and G is a ℓ -group, then every simple KG -module has the form $X \uparrow_H^G$, where X is a 1-dimensional KH -module for some subgroup $H \leq G$.

Proof: We proceed by induction on $|G|$.

If $|G| \in \{1, \ell\}$, then G is abelian and any simple KG -module S is 1-dimensional by Corollary 12.3, so $H = G$, $X = S$ and we are done.

So assume $|G| = \ell^b$ with $b \in \mathbb{Z}_{>1}$, and let S be a simple KG -module and consider the subgroup

$$U := \{g \in G \mid g \cdot x = x \ \forall x \in S\}.$$

This is obviously a normal subgroup of G since it is the kernel of the K -representation associated to S . Hence $S = \text{Inf}_{G/U}^G(T)$ for a simple $K[G/U]$ -module T . Now, if $U \neq \{1\}$, then $|G/U| < |G|$, so by the induction hypothesis there exists a subgroup $H/U \leq G/U$ and a 1-dimensional $K[H/U]$ -module Y such that $T = \text{Ind}_{H/U}^{G/U}(Y)$. But then

$$S = \text{Inf}_{G/U}^G(T) = \text{Inf}_{G/U}^G \circ \text{Ind}_{H/U}^{G/U}(Y) = \text{Ind}_H^G \circ \text{Inf}_{H/U}^H(Y),$$

so that setting $X := \text{Inf}_{H/U}^H(Y)$ yields the result. Thus we may assume $U = \{1\}$.

If G is abelian, then all simple modules are 1-dimensional, so we are done. Assume now that G is not abelian. Then G has a normal abelian subgroup A that is not central. Indeed, to construct this subgroup A , let $Z_2(G)$ denote the second centre of G , that is, the preimage in G of $Z(G/Z(G))$ (this centre is non-trivial as $G/Z(G)$ is a non-trivial ℓ -group). If $x \in Z_2(G) \setminus Z(G)$, then $A := \langle Z(G), x \rangle$ is a normal abelian subgroup not contained in $Z(G)$. Now, applying Clifford's Theorem yields:

$$S \downarrow_A^G = S_1^{a_1} \oplus \dots \oplus S_r^{a_r}$$

where $r \in \mathbb{Z}_{>0}$, S_1, \dots, S_r are non-isomorphic simple KA -modules and $S = (S_1^{a_1}) \uparrow_{H_1}^G$, where $H_1 = \text{Stab}_G(S_1^{a_1})$. We argue that $V := S_1^{a_1}$ must be a simple KH_1 -module, since if it had a proper non-trivial submodule W , then $W \uparrow_{H_1}^G$ would be a proper non-trivial submodule of S , which is simple: a contradiction. If $H_1 \neq G$ then by the induction hypothesis $V = X \uparrow_H^{H_1}$, where $H \leq H_1$ and X is a 1-dimensional KH -module. Therefore, by transitivity of the induction, we have

$$S = (S_1^{a_1}) \uparrow_{H_1}^G = (X \uparrow_H^{H_1}) \uparrow_{H_1}^G = X \uparrow_H^G,$$

as required.

Finally, the case $H_1 = G$ cannot happen. For if it were to happen then

$$S \downarrow_A^G = S \downarrow_A^{H_1} = S_1^{a_1},$$

is simple by the weak form of Clifford's Theorem, hence of dimension 1 since A is abelian. The elements of A must therefore act via scalar multiplication on S . Since such an action would commute with the action of G , which is faithful on S , we deduce that $A \subseteq Z(G)$, which contradicts the construction of A . ■

Remark 20.5

This result is extremely useful, for example, to construct the complex character table of an ℓ -group ($\ell \in \mathbb{P}$). Indeed, it says that we need look no further than induced linear characters. In general, a KG -module of the form $N \uparrow_H^G$ for some subgroup $H \leq G$ and some 1-dimensional KH -module is called **monomial**. A group all of whose simple $\mathbb{C}G$ -modules are monomial is called an **M -group**. (By the above ℓ -groups are M -groups.)