

Notation. Below R denotes an arbitrary unital and associative ring.

G Existence and uniqueness of composition series

One of the most important classes of modules is the class of simple modules. They can be used to analyse the structure of an arbitrary module, provided this module possesses a composition series. We recall here conditions of existence and uniqueness of composition series.

Definition G.1 (*Simple module*)

An R -module M is called **simple** (or **irreducible**) if it has exactly two R -submodules, namely M itself and the zero submodule.

Definition G.2 (*Composition series / composition factors / composition length*)

Let M be an R -module.

(a) A **series** (or **filtration**) of M is a finite chain of submodules

$$0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_n = M \quad (n \in \mathbb{Z}_{\geq 0}).$$

(b) A **composition series** of M is a series

$$0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_n = M \quad (n \in \mathbb{Z}_{\geq 0})$$

where M_i/M_{i-1} is simple for each $1 \leq i \leq n$. The quotient modules M_i/M_{i-1} are called the **composition factors** (or the **constituents**) of M and the integer n is called the **composition length** of M .

Clearly, in a composition series all inclusions are in fact strict because the quotient modules are required to be simple, hence non-zero.

The existence of a *composition series* implies that the module is *finitely generated*. However, the converse does not hold in general. This is explained through the fact that the existence of a composition series is equivalent to the fact that the module is both *Noetherian* and *Artinian*.

Definition G.3 (Chain conditions / Artinian and Noetherian rings and modules)

- (a) An R -module M is said to satisfy the **descending chain condition** (D.C.C.) on submodules (or to be **Artinian**) if every descending chain $M = M_0 \supseteq M_1 \supseteq \dots \supseteq M_r \supseteq \dots \supseteq \{0\}$ of submodules eventually becomes stationary, i.e. $\exists m_0$ such that $M_m = M_{m_0}$ for every $m \geq m_0$.
- (b) An R -module M is said to satisfy the **ascending chain condition** (A.C.C.) on submodules (or to be **Noetherian**) if every ascending chain $0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_r \subseteq \dots \subseteq M$ of submodules eventually becomes stationary, i.e. $\exists m_0$ such that $M_m = M_{m_0}$ for every $m \geq m_0$.
- (c) The ring R is called **left Artinian** (resp. **left Noetherian**) if the regular module R° (i.e. R itself seen as a left R -module over itself via left multiplication) is Artinian (resp. Noetherian).

Theorem G.4 (Jordan-Hölder)

Any series of R -submodules $0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_r = M$ ($r \in \mathbb{Z}_{\geq 0}$) of an R -module M may be refined to a composition series of M . In addition, if

$$0 = M_0 \subsetneq M_1 \subsetneq \dots \subsetneq M_n = M \quad (n \in \mathbb{Z}_{\geq 0})$$

and

$$0 = M'_0 \subsetneq M'_1 \subsetneq \dots \subsetneq M'_m = M \quad (m \in \mathbb{Z}_{\geq 0})$$

are two composition series of M , then $m = n$ and there exists a permutation $\pi \in \mathfrak{S}_n$ such that $M'_i/M'_{i-1} \cong M_{\pi(i)}/M_{\pi(i)-1}$ for every $1 \leq i \leq n$.

In particular, the composition length is well-defined.

Corollary G.5

If M is an R -module, then TFAE:

- (a) M has a composition series;
- (b) M satisfies D.C.C. and A.C.C. on submodules;
- (c) M satisfies D.C.C. on submodules and every submodule of M is finitely generated.

Theorem G.6 (Hopkins' Theorem)

If M is a module over a left Artinian ring R , then TFAE:

- (a) M has a composition series;
- (b) M satisfies D.C.C. on submodules;
- (c) M satisfies A.C.C. on submodules;
- (d) M is finitely generated.